Math 105 — Second Midterm March 20, 2012

Name: ____

Instructor: ____


Section: _

1. Do not open this exam until you are told to do so.

- 2. This exam has 10 pages including this cover. There are 10 problems. Note that the problems are not of equal difficulty, so you may want to skip over and return to a problem on which you are stuck.
- 3. Do not separate the pages of this exam. If they do become separated, write your name on every page and point this out to your instructor when you hand in the exam.
- 4. Please read the instructions for each individual problem carefully. One of the skills being tested on this exam is your ability to interpret mathematical questions, so instructors will not answer questions about exam problems during the exam.
- 5. Show an appropriate amount of work (including appropriate explanation) for each problem, so that graders can see not only your answer but how you obtained it. Include units in your answer where that is appropriate.
- 6. You may use any calculator except a TI-92 (or other calculator with a full alphanumeric keypad). However, you must show work for any calculation which we have learned how to do in this course.
- 7. If you use graphs or tables to find an answer, be sure to include an explanation and sketch of the graph, and to write out the entries of the table that you use.
- 8. Turn off all cell phones and pagers, and remove all headphones.
- 9. You must use the methods learned in this course to solve all problems.

Problem	Points	Score
1	14	
2	12	
3	5	
4	7	
5	12	
6	13	
7	10	
8	8	
9	9	
10	10	
Total	100	

1. [14 points] Note: No work or explanation is required on this page. The graph of a sinusoidal function g is shown below.



b. [8 points] Below are the graphs of several transformations of g(x). For each of these graphs, write the letter of the ONE function from the "Answer Choices" whose graph is shown. (**Clearly** write the capital letter of your choice on the answer blank provided.)

	Answer Cho	vices	
C. $\frac{1}{2}g(x)$ G.	$g(2x) \ g(rac{1}{2}x) \ g(x) - 1 \ g(x+1)$	I. $g(x-1)$ J. $g(x) + 1$ K. $g(x) - 2$ L. $g(x+2)$	N. $g(x) + 2$
Answer: $2 + 2$		2 + 2	Answer:
	$\begin{array}{c} & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \end{array}$		$\begin{array}{c c} & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\$
$\begin{array}{c} y \\ \hline \\ -4 \\ \hline \\ -2 \\ \hline \\ -2 \\ \hline \end{array}$		y 2 -1 -2 -2	x
Answer: †		Ť	Answer:

Show your work step-by-step and give the **exact solutions** in the answer blanks provided. **a.** [3 points] $4(3.2)^t = 7$

Answer: *t* = _____

b. [3 points] $3e^{\ln(x+2)} = 8$

Answer: *x* = _____

c. [3 points] $e^{m+5} = 6e^{-3m}$

Answer: *m* = _____

d. [3 points] $\ln(y+3) - \ln(1-y) = \ln(6)$

Answer: *y* = _____

3. [5 points] A mysterious substance decays by 30% every 6 years. Find the half-life of this substance. (Show your work carefully and either give your answer in exact form or round your answer to the nearest 0.01 year.)

Answer:

4. [7 points] Consider the function B defined by B(x) = 15 - e^{-0.001x}.
a. [3 points] Let f(x) = e^x. Use transformations to find a formula for B(x) in terms of f.


 $B(x) = _$

b. [4 points] Find the vertical and horizontal asymptotes of the graph of y = B(x). (If there are no vertical or no horizontal asymptotes, write "NONE" on the appropriate line(s).)

Vertical asymptote(s):

Horizontal asymptote(s):

- 5. [12 points] Note: You do not have to show any work on this page.
 - **a**. [6 points] If (2, -6) is a point on the graph of y = h(x), find a point on the graph of each of the functions below.



b. [6 points] Some data for functions g and k is provided in the table below. Use this data to answer the questions that follow.

x	1	2	3
g(x)	4	-1	-2
k(x)	5	4	1

(i) If g(x) is an even function, find g(-2).

Answer: g(-2) =_____

(ii) Let m(t) = 2k(-t+1). Find m(-2).

Answer: m(-2) = _____

(iii) Let n(x) = k(x-1). If n(x) is an odd function, find k(-3).

Answer: k(-3) = _____

6. [13 points] A study of mammals in a particular county in Michigan found that at the time of the study there were N groundhogs and that the population of groundhogs was increasing at a rate of 5% per year. Let G(t) be the number of groundhogs in the county t years after the study.

For full credit on this problem, you must solve for all answers algebraically and show all work step-by-step. Answers should either be in exact form or be given to at least four decimal places. **a.** [2 points] Find a formula for G(t).

Answer: G(t) =_____.

b. [3 points] Find the *continuous* growth rate of the groundhog population.

Answer: _____

c. [3 points] How long will it take for the number of groundhogs to double?

Answer: _____


d. [5 points] In the same study, it was determined that the number of moles and rabbits in the county t years after the study would be given by the formulas $M(t) = 500(0.99)^t$ and $R(t) = 200e^{0.1t}$, respectively. According to these models, when will the population of rabbits be 50% larger than the population of moles?


Answer: ____

7. [10 points] A "Whirlydoodle"¹ is a small windmill that spins and lights up when the wind blows. One evening, there is a light breeze and a particular Whirlydoodle's blades are rotating at a constant rate of one revolution every 4 seconds. A moth lands on the tip of one of the blades of the Whirlydoodle when the blade is pointed straight up. (The moth then hangs on and rides for a minute.) This Whirlydoodle is mounted 5 feet (60 inches) above the ground, and each blade is 10 inches long, as shown in the diagram on the right.

Let h(t) be the height (in inches) of the moth above the ground t seconds after the moth lands on the Whirlydoodle.

a. [6 points] Sketch a graph of y = h(t) for $0 \le t \le 8$. (Remember to label the axes (including units) and to make sure that the key features and characteristics of your graph are clear.)





b. [4 points] Find a formula for h(t).

Answer: h(t) =_

 $^{^1}$ "Whirly doodles" can be seen around downtown Ann Arbor.

8. [8 points] Throughout this problem, make sure that you clearly show your work step-by-step. In the 1970's, seismologists developed the Moment Magnitude Scale (MMS) to estimate the magnitude of large earthquakes in terms of the energy released. Unlike the Richter scale, which is based on the size of seismic waves, the MMS is based on seismic moments (which represent the energy released in an earthquake). The MMS rating of an earthquake is defined to be

$$S = \frac{2}{3} \log\left(\frac{M}{A}\right)$$

where M is the seismic moment of the quake (in dynes/cm) and A is a positive constant.


a. [4 points] Let S_1 and S_2 represent the MMS ratings of two earthquakes with seismic moments M_1 and M_2 , respectively. Using properties of logarithms, find a formula for $S_2 - S_1$ in terms of M_1 and M_2 . Simplify your formula as much as possible.

Answer: $S_2 - S_1 =$ _____.

b. [4 points] The San Francisco earthquake of 1989 had an MMS rating of 6.9 and the Northridge, CA earthquake of 1994 had an MMS rating of 6.7. Based on these ratings, how many times greater than the Northridge seismic moment was the San Francisco seismic moment? (Give your answer in exact form or round to the nearest 0.01.)

Answer: The seismic moment from the San Francisco earthquake was ______ times greater than the seismic moment of the Northridge earthquake.

9. [9 points] Consider the points P and Q determined by the angle $\frac{2\pi}{7}$ as shown in the diagram below.



You don't have to show work, but any work you do show may be considered for partial credit. Give all answers in *exact form*.

a. [2 points] Find the coordinates of the point P.

Answer: The coordinates of P are (______, ____).

b. [2 points] Find the coordinates of the point Q.

Answer: The coordinates of Q are (______, ____).

c. [2 points] Find the length of the counterclockwise path from the point Q to the point (-1.5, 0). (This path is shown in **bold** in the diagram above.)

Answer: _____

d. [3 points] An ant begins at the point P, walks *clockwise* along the unit circle for 3 units and then stops. What are the coordinates of the point at which the ant stops?

Answer: The coordinates of this point are (______, ____)

10. [10 points] Ivanka is a student at a nearby college. Let C(h) be the total tuition, in thousands of dollars, the college charges her if she takes h credit hours, and let a be the average number of credit hours students take at the college.

For each of the following, pick the ONE expression from the list of "Answer Choices" that best represents the described quantity. Clearly write the capital letter of your choice on the answer blank provided.

Answer Choices							
А.	C(3)	F.	C(a) + 3	К.	$3C^{-1}(a)$	Р.	$C^{-1}(3a)$
В.	$C^{-1}(3)$	G.	C(a-3)	L.	$C^{-1}(a)/3$	Q.	3C(C(a))
С.	C(a)	Η.	C(a) - 3	М.	C(a)/3	R.	$C(3C^{-1}(a))$
D.	$C^{-1}(a)$	I.	3C(a)	Ν.	C(a/3)	S.	$C^{-1}(3C(a))$
Е.	C(a+3)	J.	C(3a)	О.	$C^{-1}(a/3)$	Т.	$C^{-1}(C(3a))$

- **a**. [2 points] Ivanka's tuition (in thousands of dollars) if she takes a total of 3 credit hours
- **b**. [2 points] Ivanka's total tuition (in thousands of dollars) if she takes 3 credit hours more than average
- c. [2 points] Ivanka's tuition (in thousands of dollars) if she takes one third the average number of credit hours

Answer:

d. [2 points] The amount (in thousands of dollars) that Ivanka pays for tuition if she takes the average number of credit hours but has a scholarship that covers three thousand dollars of her tuition

Answer: _____

e. [2 points] The number of credit hours Ivanka takes if her total tuition is three times as much as the tuition for taking the average number of credit hours

Answer: _____

Answer: _____

Answer: _____