


1. [10 points] Given below are three functions.  $r(w)$  is given by a graph,  $h(t)$  is given by a formula, and  $n(v)$  is described verbally.

$n(v)$  has a constant rate of change, and its graph passes through the points  $(1, 4)$  and  $(3, 0)$ .

$$h(t) = \sqrt{t - 4}.$$



The function  $r(w)$  is linear on  $[-2, 0]$  and on  $[2, 4]$ . Give your answer in **exact** form (i.e. no decimal approximations) for parts **a.-c.**

- a. [2 points] Complete the sentence by filling in the blank. You can express your answer in inequality or interval notation.

The domain of  $h(t)$  is \_\_\_\_\_.

- b. [2 points] Complete the sentence by filling in the blank. You can express your answer in inequality or interval notation.

The range of  $r(w)$  is \_\_\_\_\_.

- c. [2 points] Complete the sentence by filling in the blank.

The average rate of change of  $h(t)$  between  $t = 6$  and  $t = 9$  is \_\_\_\_\_.

- d. [4 points] Find all solutions to the equation

$$n(r(w)) = -2.$$

If there is no solution, write “no solution” in the blank. Show your work. (If needed, use the graph of  $r(w)$  to give estimates for values of  $w$  in the interval  $[0, 2]$ . Otherwise, give your answer in exact form.)

$w =$  \_\_\_\_\_.