

1. [5 points] For each of the statements below, circle “**True**” if the statement is *definitely* true. Otherwise, circle “**False**”. You do not need to show any work for this problem.

- a. [1 point] If a function has more than one zero, then the function is not invertible. True False
- b. [1 point] If $x > 1$, then $100x^{100000} > e^{0.0001x}$. True False
- c. [1 point] If $h(t) = \ln(t)$ then $h^{-1}(t) = \frac{1}{\ln(t)}$. True False
- d. [1 point] If a function is concave up, then the function is increasing. True False
- e. [1 point] If $f(x)$ and $g(x)$ are both even functions, then the function $f(g(x))$ is also an even function. True False

2. [6 points] Solve each of the equations below. *Show your work step-by-step and write the solutions in exact form in the answer blanks provided.*

a. [3 points] $5e^{2t+7} = 3(4^t)$

Solution: We first divide both sides of this equation by 5 to find $e^{2t+7} = 0.6(4^t)$. Then we use logarithms to find t .

$$\begin{aligned}\ln(e^{2t+7}) &= \ln(0.6(4^t)) \\ 2t + 7 &= \ln(0.6) + \ln(4^t) = \ln(0.6) + t \ln(4) \\ 2t - t \ln(4) &= \ln(0.6) - 7 \\ t(2 - \ln(4)) &= \ln(0.6) - 7 \text{ so } t = \frac{\ln(0.6) - 7}{2 - \ln(4)}\end{aligned}$$

Answer: $t = \frac{\ln(0.6) - 7}{2 - \ln(4)}$

b. [3 points] $\log(w) + \log(w + 3) = 1$

Solution: We apply a basic property of logarithms and then use the definition or the logarithm (or exponentiate) to solve for w .

$$\begin{aligned}\log(w) + \log(w + 3) &= 1 \\ \log(w(w + 3)) &= 1 \\ w(w + 3) &= 10^1 \\ w^2 + 3w &= 10 \\ w^2 + 3w - 10 &= 0 \\ (w - 5)(w + 2) &= 0 \\ w &= 5 \text{ or } w = -2\end{aligned}$$

However, note that $w = -2$ is not a solution to the original equation because -2 is not in the domain of $\log w$. Hence the only solution is $w = 5$.

Answer: $w = 5$