

MATH 115 – SECOND MIDTERM EXAM

March 27, 2007

NAME: _____

INSTRUCTOR: _____ SECTION NUMBER: _____

1. **Do not open this exam until you are told to begin.**
2. This exam has 9 pages including this cover. There are 8 questions.
3. Do not separate the pages of the exam. If any pages do become separated, write your name on them and point them out to your instructor when you turn in the exam.
4. Please read the instructions for each individual exercise carefully. One of the skills being tested on this exam is your ability to interpret questions, so instructors will not answer questions about exam problems during the exam.
5. Show an appropriate amount of work for each exercise so that the graders can see not only the answer but also how you obtained it. Include units in your answers where appropriate.
6. You may use your calculator. You are also allowed two sides of a 3 by 5 notecard.
7. If you use graphs or tables to obtain an answer, be certain to provide an explanation and sketch of the graph to show how you arrived at your solution.
8. Please turn off all cell phones and pagers and remove all headphones.

PROBLEM	POINTS	SCORE
1	16	
2	16	
3	12	
4	14	
5	10	
6	12	
7	14	
8	6	
TOTAL	100	

1. (4 points each) For the following statements circle True or False. If the statement is *always* true, explain why it is true. If it is false give an example of when the statement is false. Examples may be formulas or graphs.

(a) If $y(x)$ is a twice differentiable function, then $\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$.

True

False

- (b) There exists a function $f(x)$ such that $f(x) > 0$, $f'(x) < 0$, and $f''(x) > 0$ for all real values of x .

True

False

- (c) If h is differentiable for all x and $h'(a) = 0$, then $h(x)$ has a local minimum or local maximum at $x = a$.

True

False

- (d) If f and g are positive and increasing on an interval I , then f times g is increasing on I .

True

False

2. (4 points each) Suppose f and g are differentiable functions with values given by the table below:

x	$f(x)$	$g(x)$	$f'(x)$	$g'(x)$
1	2	9	-3	7
3	4	11	15	-19


(a) If $h(x) = f(x)g(x)$, find $h'(3)$.

(b) If $j(x) = \frac{(g(x))^3}{f(x)}$, find $j'(1)$.

(c) If $d(x) = x \ln(e^{f(x)})$, find $d'(3)$.

(d) If $t(x) = \cos(g(x))$, find $t'(1)$.

3. (6 points each) Consider the graphs of $m(x)$ and $n(x)$ below. Let $h(x) = n(m(x))$. Find the following, or explain why they do not exist. The function m has a sharp corner at $x = 50$ and n has a sharp corner at $x = 80$. Determine values that exist as *exact* values—*i.e.*, not a graphical approximation. Please circle your answers.



(a) $h'(80)$

(b) a value of x such that $h'(x) = -2$

4. Suppose that x and y satisfy the relation given by the curve

$$x^4 + y^3 = 2 + \frac{7}{2}xy$$

(a) (5 points) Find $\frac{dy}{dx}$.

(b) (3 points) Under what condition(s) (if any) on x and y is the tangent line to the curve horizontal?

(c) (2 points) Consider the points $(1, 2)$ and $(3, 4)$. One of these points lies on the curve, and one does not. Show which point lies on the curve and which does not.

(d) (4 points) Find an equation of the tangent line to the curve at the point from part (c) that is on the curve.

5. (10 points) Find the quadratic polynomial $g(x) = ax^2 + bx + c$ which “best fits” the function $f(x) = \ln(x)$ at $x = 1$ in the sense that

$$g(1) = f(1), \quad \text{and} \quad g'(1) = f'(1), \quad \text{and} \quad g''(1) = f''(1).$$

$$g(x) = \underline{\hspace{10cm}}$$

6. (12 points) It's time to redesign the layout of exhibits at the San Diego Zoo. The zookeeper, Joan Embery, has been told by Paco Underhill that more exhibits will attract more visitors to enter the zoo but, as the space between exhibits decreases, more zoo visitors are likely to brush butts and flee the zoo in disgust.¹

Paco has modeled the predicted number of visitors to the zoo each year, in millions of people, by the function

$$f(x) = axe^{-bx} + c$$

where x represents the number of exhibits per acre.

Since the park is beautiful on its own, Paco believes that 1/2 million visitors a year will come to the area, even if there are no exhibits. He has determined that the maximum number of visitors will come to the zoo if there are 4 exhibits per acre. (After that, the “disgust factor” begins to creep in.) According to Paco’s model, when the number of exhibits per acre is 4, the number of visitors would be approximately 2.5 million people per year. Use this information (and calculus) to solve for a , b and c in the above function.

[Note: The maximum number of exhibits that could be packed into an acre of land is 8, since exhibits require 500 m², on average, and an acre \approx 4000 m².]


$$a = \underline{\hspace{2cm}}$$

$$b = \underline{\hspace{2cm}}$$

$$c = \underline{\hspace{2cm}}$$

¹See http://en.wikipedia.org/wiki/Joan_Embery. and <http://www.amazon.com/Why-We-Buy-Science-Shopping/dp/0684849143>. When he finishes with the zoo, Underhill will arrange the classroom tables in Dennison Hall.

7. (14 points) No matter what is done with the other exhibits, the octopus tank at the zoo must be rebuilt. (The current tank has safety issues, and there are fears that the giant octopus might escape!) The new tank will be 10 feet high and box-shaped. It will have a front made out of glass. The back, floor, and two sides will be made out of concrete, and there will be no top. The tank must contain at least 1000 cubic feet of water. If concrete walls cost \$2 per square foot and glass costs \$10 per square foot, use calculus to find the dimensions and cost of the least-expensive new tank. [Be sure to show all work.]



GIANT OCTOPUS (*Enteroctopus*)²

Dimensions: _____

Minimum Cost: _____

²See <http://www.cephbase.utmb.edu/Tcp/pdf/anderson-wood.pdf>. (They really DO escape....)

8. (6 points) On the axes below, sketch a possible graph of a single function $y = g(x)$ satisfying all of the properties below: [Label your points on the axes.]

(i) $g(x)$ is defined and continuous for all values of x .

(ii) $g(x)$ has critical points at $x = -1$ and $x = 4$.

(iii) $g'(x) \geq 0$ on $(-\infty, 4)$.

(iv) $g(x)$ is decreasing on $(4, \infty)$.

(v) $\lim_{x \rightarrow \infty} g(x) = -2$.

