7. [12 points] The graph of the function f(x) is shown below.



For a.-c., give your answers as a list of one or more of the given numbers, or write NONE.

- **a**. [1 point] At which of the values a = 1, 2, 3, 4, 5 is f(a) undefined?
- **b.** [1 point] For which of the values a = 1, 2, 3, 4, 5 is f(x) continuous at x = a?
- **c**. [2 points] For which of the values a = 1, 2, 3, 4, 5 is $f(a) = \lim_{x \to a^-} f(x)$?

For d.-g., use the graph of the function f(x) to evaluate each of the expressions below. If a limit diverges to ∞ or $-\infty$ or if the limit does not exist for any other reason, write DNE.

d. [2 points] $\lim_{x \to 5} f(x)$ **e.** [2 points] $\lim_{x \to 3} f(x)$ **f.** [2 points] $\lim_{x \to 0} f(4 + |x|)$ **g.** [2 points] $\lim_{h \to 0} \frac{f(4.25 + h) - f(4.25)}{h}$

8. [11 points] Consider the rational function $g(x) = \frac{(x-12)(x-7)(x-2)}{(2x-4)(x-3)(x-5)}$.

- **a**. [2 points] What are the vertical asymptotes of the function g(x)?
- **b.** [2 points] What are the vertical asymptotes of the function $\frac{1}{g(x)}$?

The piecewise function h(x) is defined as follows, where g(x) is as above, where f(x) is from Problem 7 above, and where B is a nonzero constant.

$$h(x) = \begin{cases} \frac{e^{2x}}{x^2} & x \le 3\\ B \cdot f(x) & 3 < x \le 6\\ g(x) & 6 < x \end{cases}$$

c. [3 points] Find an *exact* value of B for which the function h(x) is continuous at x = 3. Show your work.

Evaluate each of the expressions below. If a limit diverges to ∞ or $-\infty$ or if the limit does not exist for any other reason, write DNE.

d. [2 points] $\lim_{x \to \infty} h(x)$ **e.** [2 points] $\lim_{x \to -\infty} h(x)$