3. [4 points] An insect population in a certain large park varies sinusoidally from a low of 10 million on January 1st to a high of 70 million on July 1st. Let P(t) be the population in the park of this insect, in millions, t months after January 1st. Find a formula for P(t).

Solution: We are looking for a sinusoidal function, of the form $y = A \sin (B(t-h)) + C$ or $y = A \cos (B(t-h)) + C$. Since P(0) is a minimum, we will use cosine instead of sine, with A negative, and we do not need any phase shift so h = 0. Since P(t) varies between 10 and 70, its amplitude will be $\frac{70-10}{2} = 30$, and the midline will be $\frac{10+70}{2} = 40$. So A = -30 and C = 40. Finally, since the period is 12 months, we have $B = \frac{2\pi}{12}$. Therefore,

$$P(t) = -30\cos\left(\frac{2\pi t}{12}\right) + 40.$$

Answer:
$$P(t) = -30\cos(\frac{2\pi t}{12}) + 40$$

 $(2\pi t)$

4. [5 points] Let

$$g(x) = \begin{cases} \frac{\arctan x}{x} & x \neq 0, \\ 1 & x = 0. \end{cases}$$

You are given that g'(0) exists. Use the limit definition of the derivative to write an explicit expression for g'(0). Your answer should not involve the letter g. Do not attempt to evaluate or simplify the limit. Write your final answer in the answer box provided below.

Solution:

$$g'(0) = \lim_{h \to 0} \frac{g(0+h) - g(0)}{h} = \lim_{h \to 0} \frac{\frac{\arctan(0+h)}{0+h} - 1}{h} = \lim_{h \to 0} \frac{\frac{\arctan(h)}{h} - 1}{h}.$$

Answer:
$$g'(0) = \lim_{h \to 0} \frac{\frac{\arctan(h)}{h} - 1}{h}$$