

10. [7 points] The curve  $\mathcal{C}$  is given by the equation  $e^{\cos(x^2-y^2)} = ex$ .

a. [2 points] Which of the following points  $(x, y)$  lie on the curve  $\mathcal{C}$ ? Circle all correct answers.

(1, 1)      (-1, 1)       (1, -1)      (0,  $\sqrt{2}$ )      NONE OF THESE

b. [5 points] Compute  $\frac{dy}{dx}$ . Show every step of your work and circle your final answer.

*Solution:* Implicitly differentiating, we get:  $e^{\cos(x^2-y^2)} \cdot (-\sin(x^2-y^2)) \cdot \left(2x - 2y\frac{dy}{dx}\right) = e$ .

Now we solve for  $\frac{dy}{dx}$ :

$$2x - 2y\frac{dy}{dx} = \frac{e}{e^{\cos(x^2-y^2)} \cdot (-\sin(x^2-y^2))},$$

$$-2y\frac{dy}{dx} = \frac{e}{e^{\cos(x^2-y^2)} \cdot (-\sin(x^2-y^2))} - 2x,$$

$$\frac{dy}{dx} = \frac{e}{-2ye^{\cos(x^2-y^2)} \cdot (-\sin(x^2-y^2))} - \frac{2x}{-2y} = \frac{e}{2ye^{\cos(x^2-y^2)} \sin(x^2-y^2)} + \frac{x}{y}.$$

Or, one can distribute the left-hand side first, to obtain the equivalent answer

$$\frac{dy}{dx} = \frac{2xe^{\cos(x^2-y^2)} \cdot \sin(x^2-y^2) + e}{2ye^{\cos(x^2-y^2)} \cdot \sin(x^2-y^2)}.$$

11. [8 points]

Suppose  $h(x)$  is a function such that  $h(x)$  has **exactly three** critical points. Assume that both  $h(x)$  and  $h'(x)$  are differentiable on  $(-\infty, \infty)$ . A table of values is given to the right.

$x$	0	3	5	7
$h(x)$	2	?	4	4
$h'(x)$	-1	0	0	?

a. [2 points] Note that  $h(x)$  satisfies the hypotheses of the Mean Value Theorem on  $[5, 7]$ . Briefly explain why the conclusion of this theorem implies that one of the three critical points of  $h(x)$  must be in the interval  $5 < x < 7$ .

*Solution:* The conclusion of the Mean Value Theorem applied to  $h(x)$  on  $[5, 7]$  states that there exists a point  $c$  with  $5 < c < 7$  such that  $h'(c) = \frac{h(7) - h(5)}{7 - 5} = \frac{4 - 4}{7 - 5} = 0$ , making  $c$  a critical point of  $h(x)$  in the interval  $5 < x < 7$ .

In the parts below, circle all correct answers. No justification is needed.

b. [2 points] On which of the following intervals *must*  $h(x)$  be increasing on the entire interval?

(0, 3)       (3, 5)      (5, 6)      (6, 7)      NONE OF THESE

c. [2 points] Which of the following *could* be the  $x$ -coordinate of a global minimum of  $h(x)$  on  $(-\infty, \infty)$ ?

$x = 0$         $x = 3$        $x = 5$         $x = 6$       NONE OF THESE

d. [2 points] Also suppose that  $h(x)$  is concave down on  $(-\infty, 0)$ . Which of the following *could* be the value of  $h(-2)$ ?

1      2       3      4      5      NONE OF THESE