

8. [6 points] The equation $x^2 + xy + 2y^2 = 28$ defines y implicitly as a function of x .

a. [4 points] Compute $\frac{dy}{dx}$. Show every step of your work.

Answer: _____

b. [2 points] Find an equation of the line tangent to the curve defined by $x^2 + xy + 2y^2 = 28$ at the point $(2, 3)$.

Answer: _____

9. [6 points] The equation $x + \frac{1}{3}y^3 - y = 1$ implicitly defines x and y as functions of each other. Implicitly differentiating this equation with respect to x and solving for $\frac{dy}{dx}$ gives

$$\frac{dy}{dx} = \frac{-1}{y^2 - 1}.$$

Let \mathcal{C} be the graph of the equation $x + \frac{1}{3}y^3 - y = 1$. Note that all points listed as possible answers below do actually lie on the graph \mathcal{C} .

a. [2 points] Circle all points below at which the line tangent to \mathcal{C} is *horizontal*.

$(-5, 3)$ $(\frac{1}{3}, -1)$ $(1, 0)$ $(1 + \frac{\sqrt{2}}{3}, \sqrt{2})$ $(\frac{5}{3}, 1)$ NONE OF THESE

b. [2 points] Circle all points below at which the line tangent to \mathcal{C} is *vertical*.

$(-5, 3)$ $(\frac{1}{3}, -1)$ $(1, 0)$ $(1 + \frac{\sqrt{2}}{3}, \sqrt{2})$ $(\frac{5}{3}, 1)$ NONE OF THESE

c. [2 points] Circle all points below at which $\frac{dy}{dx}$ and $\frac{dx}{dy}$ are equal to each other.

$(-5, 3)$ $(\frac{1}{3}, -1)$ $(1, 0)$ $(1 + \frac{\sqrt{2}}{3}, \sqrt{2})$ $(\frac{5}{3}, 1)$ NONE OF THESE