


10. [5 points] Shown on the axes below are the graphs of $y = f(x)$, $y = f'(x)$, and $y = f''(x)$.



Determine which graph is which and circle the ONE correct response below.

- $f(x)$: I, $f'(x)$: II, and $f''(x)$: III
- $f(x)$: I, $f'(x)$: III, and $f''(x)$: II
- $f(x)$: II, $f'(x)$: I, and $f''(x)$: III
- $f(x)$: II, $f'(x)$: III, and $f''(x)$: I
- $f(x)$: III, $f'(x)$: I, and $f''(x)$: II
- $f(x)$: III, $f'(x)$: II, and $f''(x)$: I

11. [4 points] Suppose w and r are continuous functions on $(-\infty, \infty)$, $W(x)$ is an invertible antiderivative of $w(x)$, and $R(x)$ is an antiderivative of $r(x)$.

Circle all of the statements I-VI below that must be true.

If none of the statements must be true, circle NONE OF THESE.

- I. $W(x) + R(x) + 2$ is an antiderivative of $w(x) + r(x)$.
- II. $W(x) + R(x)$ is an antiderivative of $w(x) + r(x) + 2$.
- III. $\cos(W(x))$ is an antiderivative of $\sin(w(x))$.
- IV. $e^{W(x)}$ is an antiderivative of $w(x)e^{w(x)}$.
- V. $e^{R(x)}$ is an antiderivative of $r(x)e^{R(x)}$.
- VI. If w is never zero, then $W^{-1}(R(x))$ is an antiderivative of $\frac{r(x)}{w(W^{-1}(R(x)))}$.
- VII. NONE OF THESE