

# MATH 116 — FINAL EXAM

Winter 2004

NAME: \_\_\_\_\_

ID NUMBER: \_\_\_\_\_

INSTRUCTOR: \_\_\_\_\_


SECTION NO: \_\_\_\_\_

1. Do not open this exam until you are told to begin.
2. This exam has 10 pages including this cover. There are 10 questions.
3. Do not separate the pages of the exam. If any pages do become separated, write your name on them and point them out to your instructor when you turn in the exam.
4. Please read the instructions for each individual exercise carefully. One of the skills being tested on this exam is your ability to interpret questions, so instructors will not answer questions about exam problems during the exam.
5. Show an appropriate amount of work for each exercise so that the graders can see not only the answer but also how you obtained it. Include units in your answers where appropriate.
6. You may use your calculator. You are also allowed 2 sides of a 3 by 5 notecard.
7. If you use graphs or tables to obtain an answer, be certain to provide an explanation and sketch of the graph to make clear how you arrived at your solution.
8. Please turn **off** all cell phones.

PROBLEM	POINTS	SCORE
1	4	
2	6	
3	6	
4	13	
5	10	
6	15	
7	12	
8	10	
9	10	
10	14	
TOTAL	100	

1. (4 points) Circle the differential equation whose slope field is shown in the figure.

- A.  $\frac{dy}{dx} = \sin x$     B.  $\frac{dy}{dx} = -y$     C.  $\frac{dy}{dx} = x^2 + y^2$   
 D.  $\frac{dy}{dx} = x + y$     E.  $\frac{dy}{dx} = x - 2y$     F.  $\frac{dy}{dx} = \sin(x + y)$



2. (6 points) The function  $f$  is a continuous function, some of whose values are given in the following table.

$x$	0	1	2	3	4	5	6
$f(x)$	8	6	3	-2	0	1	2

For the function  $F$  defined by  $F(x) = \int_0^x f(t)e^{-t} dt$ , what is  $F'(2)$ ?

$F'(2) = \underline{\hspace{2cm}}$ .

3. (6 points) Does the infinite series  $\sum_{n=1}^{\infty} ne^{-n^2}$  converge or diverge? (Show your work.)

4. (13 points) Consider the initial value problem

$$\frac{dy}{dx} = y + e^x, \quad y(0) = 0.$$

(a) Show that  $y = xe^x$  is a solution to the initial value problem.


(b) Compute the approximation to the solution,  $y(1)$ , of the problem at  $x = 1$  given by Euler's method with four steps (i.e.  $\Delta x = \frac{1}{4}$ ). You must show all your work to receive credit.

$y(1) \approx$  \_\_\_\_\_

(c) What is the error when the exact solution  $y(1)$  is approximated by Euler's method with 4 steps? If instead you were to use Euler's method with 16 steps, approximately what would you expect the error to be? (No explanation required. DO NOT carry out Euler's method with 16 steps.)

4-step error = \_\_\_\_\_ 16-step error estimate  $\approx$  \_\_\_\_\_

5. (10 points) The number of micrograms  $x$  of pollen produced annually by plants of a certain species in a small forest has been determined (by experiment) to have a density function  $p(x)$  whose graph is shown in the figure.



(a) Write a definite integral whose value is the fraction of the plants that produce less than 10 micrograms of pollen each year.

(b) What fraction of the plant population produces less than 10 micrograms of pollen each year?

(c) Let  $P(x)$  be the cumulative distribution function for this population. In terms of the population, what is the meaning of  $P(13) - P(8)$ ?

(d) What is the median number of micrograms of pollen produced by plants in this population?

6. (15 points) For each of the following statements, circle **T** if the statement is always true, and otherwise circle **F**. No explanations are required.

(a) The Taylor series for  $\sin(x)$  about  $x = 1$  is  $(x - 1) - \frac{(x - 1)^3}{3!} + \frac{(x - 1)^5}{5!} - \dots$

**T**                      **F**

(b) If Euler's method with 10 steps is used to approximate the solution to the initial value problem  $\frac{dy}{dx} = -y$ ,  $y(0) = 1$  at  $x = 1$ , then the approximation will be an overestimate for the exact solution.

**T**                      **F**

(c) Let  $f$  be a continuous, positive, decreasing function defined for  $x \geq 1$  such that  $\int_1^{\infty} f(x) dx$  converges. If  $a_n = f(n)$ , then  $\sum_{n=1}^{\infty} a_n = \int_1^{\infty} f(x) dx$ .

**T**                      **F**

(d) The system of differential equations,

$$\begin{aligned} \frac{1}{x} \frac{dx}{dt} &= y - 1, \\ \frac{1}{y} \frac{dy}{dt} &= x - 1, \end{aligned}$$

models the interaction of two populations involved in a predator-prey relationship.

**T**                      **F**

(e) The relative growth rate of the population in the *logistic model for population growth* is a linear function of the population.

**T**                      **F**

**7.** (12 points) A hard-boiled egg at  $98^{\circ}\text{C}$  is put in a sink of  $18^{\circ}\text{C}$  water. After 5 minutes, the egg's temperature is  $38^{\circ}\text{C}$ . Assume that the water has not warmed appreciably and that the temperature of the egg changes at a rate proportional to the difference between its temperature and that of the water.

**(a)** Write the differential equation and initial conditions which model the temperature of the egg.

**(b)** Find the solution of the initial value problem of part (a). (Show your work.)

**(c)** How long does it take for the temperature of the egg to reach  $20^{\circ}\text{C}$ ?

**8.** (10 points) The *electric potential* is a quantity of great importance in electrostatics. The electric potential  $V(R)$  at a distance  $R$  along the axis perpendicular to the center of a charged disk with radius 1 is given by

$$V(R) = C \left( \sqrt{R^2 + 1} - R \right)$$

where  $C$  is a constant that depends on the choice of units that are being used.

(a) Show that for large numbers  $R$ ,

$$V(R) \approx \frac{C}{2R}.$$

(Hint:  $\sqrt{R^2 + 1} = R\sqrt{1 + \frac{1}{R^2}}$  and remember that  $R$  is large.)

(b) Approximately how large should  $R$  be in order that the error in the approximation of  $V(R)$  by  $C/2R$  is less than 4% of  $V(R)$ ?

**9.** (10 points). The watering tank for livestock on a farm has depth of 30 inches (and a constant cross-sectional area). The tank is connected to a water source and controller so that when the depth of water in the tank is less than 25 inches, water flows into the tank at a constant rate that increases the depth of water by 4 inches per hour. When the depth of water is greater than 25 inches, no water flows into the tank.

At 6 am one morning, when the depth of the water in the tank is 25 inches, it springs a leak and the water leaks from the tank at a rate proportional to the square root of the depth of water. Let  $c$  be the proportionality constant.

**(a)** Write the differential equation for the depth  $h(t)$  of the water in the tank at time  $t$ , where  $t$  is the time in hours after 6 am. Assume that when there is 25 inches of water in the tank, the proportionality constant  $c$  is large enough that the water leaks out faster than the four inch per hour rate at which the source adds water. (Do not attempt to solve the differential equation.)

**(b)** Unfortunately, the farmer falls ill and no one checks on the tank to discover the leak for a very long time. When the leak is finally discovered, it is also found that the depth of water in the tank is 9 inches. Based on this information, estimate the constant of proportionality,  $c$ , that determines the rate that water leaks from the tank? Explain how you found your answer.



**10.** (14 points) A cylindrical tank is has a circular cross section of radius 2 meters and a length of 4 meters. It is to be filled with a compressible liquid whose density varies with its height and is equal to  $\rho(h) = 60\sqrt{1+h}$  kg/m<sup>3</sup> at  $h$  meters below the surface of the liquid.

**(a)** Suppose the tank is standing on one of its circular ends (figure 1) and is filled with the liquid. What is the approximate mass in a thin horizontal slice of thickness  $\Delta h$  that is  $h$  meters below the top of the tank?




Figure 1

**(b)** Write a definite integral whose value is equal to the total mass of the liquid in the tank.

**(c)** Evaluate your integral from part (b) to find the total amount of liquid in the tank. Show your work, or explain how you obtained your answer.

(d) Suppose that instead the tank is lying on a side (figure 2) and again filled with the liquid. What then is the approximate mass of the liquid in a thin horizontal slice of thickness  $\Delta h$  that is  $h$  meters below the top of the tank?




Figure 2

(e) Is the amount of liquid the same as in part (c)? If so, explain why. If not, find the amount of liquid in the tank.