

Math 116 — First Midterm

February 2011

Name: _____

Instructor: _____ Section: _____

1. **Do not open this exam until you are told to do so.**
 2. This exam has 10 pages including this cover. There are 8 problems. Note that the problems are not of equal difficulty, so you may want to skip over and return to a problem on which you are stuck.
 3. Do not separate the pages of this exam. If they do become separated, write your name on every page and point this out to your instructor when you hand in the exam.
 4. Please read the instructions for each individual problem carefully. One of the skills being tested on this exam is your ability to interpret mathematical questions, so instructors will not answer questions about exam problems during the exam.
 5. Show an appropriate amount of work (including appropriate explanation) for each problem, so that graders can see not only your answer but how you obtained it. Include units in your answer where that is appropriate.
 6. You may use any calculator except a TI-92 (or other calculator with a full alphanumeric keypad). However, you must show work for any calculation which we have learned how to do in this course. You are also allowed two sides of a $3'' \times 5''$ note card.
 7. If you use graphs or tables to find an answer, be sure to include an explanation and sketch of the graph, and to write out the entries of the table that you use.
 8. **Turn off all cell phones and pagers**, and remove all headphones.
-

Problem	Points	Score
1	10	
2	17	
3	15	
4	15	
5	8	
6	12	
7	11	
8	12	
Total	100	

1. [10 points] Indicate if each of the following statements are true or false by circling the correct answer. **Justify your answers.**

a. [2 points] If $F(x)$ is an antiderivative of an even function $f(x)$, then $F(x)$ must also be an even function.

True False

b. [2 points] If $G(x)$ is an antiderivative of $g(x)$ and $(G(x) - F(x))' = 0$, then $F(x)$ is an antiderivative of $g(x)$.

True False

c. [2 points] Let $f(t) = bt + ct^2$ with $b > 0$ and $c > 0$, then $\text{Left}(n) \leq \int_0^{10} f(t)dt$ for all n .


True False

d. [2 points] The average of an even function $f(x)$ over the interval $[-a, a]$ is equal to twice its average over the interval $[0, a]$.

True False


e. [2 points] The density δ of a circular porcelain dinner plate depends on the distance r from the center of the plate. The relationship between δ and r is shown in the graph below. The center of mass of this plate is located near the edge of the plate.

True False



2. [17 points]


The graph of an odd function f is shown below.



- a. [7 points] Let $F(x)$ be the antiderivative of $f(x)$ with the property that $F(3) = -2$. Use the graph of $f(x)$ to compute the following values of $F(x)$.



x	-7	-6	-3	0	3	6	7
$F(x)$							

- b. [8 points] Sketch the graph of $F(x)$ from $x = -7$ to $x = 7$. Label all points of inflection.



- c. [2 points] Calculate the average value of f between $x = -3$ and $x = 7$.

3. [15 points] Use the graphs of $f(x)$ and $g(x)$ to find the EXACT values of $A, B,$ and C . Show all your work.



a. [2 points] $A = \int_{-3}^6 |f(x)| dx$

b. [5 points] $B = \int_0^2 xg'(x^2) dx$

c. [8 points] $C = \int_0^3 2xg'(x) dx$

4. [15 points] A patient is given 100 mg of an experimental drug. It has been estimated that the rate $f(t)$ at which his body eliminates the drug is given in the following table. Values of t are in hours after the administration of the drug and $f(t)$ is measured in mg/hour.

t	0	0.5	1.0	1.5	2
$f(t)$	17.3	14.5	12.2	10.3	8.6


Assume $f(t)$ is continuous with no critical points or points of inflection in $0 \leq t \leq 2$. Make sure to include the appropriate units in your answers below.

- a. [4 points] Use each left, right, trapezoid and midpoint sums to estimate amount of drug eliminated after 2 hours. When calculating each sum, use the maximum number of subdivisions possible. Show all the terms in each sum.

- b. [4 points] Using the computations in a), what is the best overestimate you can find for the amount of drug removed from the patient's body after 2 hours? What is the best underestimate? Justify your answers.

- c. [4 points] Using left and right hand sums, how often do we have to measure $f(t)$ in $0 \leq t \leq 2$ to obtain an estimate of the amount of drug eliminated from the patient's body after 2 hours within 0.1 mg of its actual value?
- d. [3 points] Find a formula for $g(t)$, the amount of drug (in mg) left in the patient's body after t hours of being administered.


5. [8 points] A company wants to design a bicycle ramp using the shape of the graph of the function $f(x) = \frac{4}{3}x^{\frac{3}{2}}$, where x is the length in meters of the base of the ramp.



Find the length s of a ramp with base of length L . Show all your work.

6. [12 points]

The region bounded by the graph of $y = e^{0.5x}$, the line $y = 1$, and the line $x = 2$ is shown below. For each of the lines L_1 and L_2 write a definite integral that represents the volume of the solid object obtained by rotating the region around that line. You do not need to show your work or calculate the value of the integral.



a. [6 points] L_1 :

b. [6 points] L_2 :

7. [11 points]

A truck carrying a large tank of paint leaves a garage at 9AM. The tank starts to leak in such a way that x miles from the garage, the density of paint on the road is $e^{-x^2/5000}$ gallons per mile. At 10AM, a cleaning crew leaves from the same garage and follows the path of the truck, scrubbing the paint from the road as it travels until it catches up to the leaking truck. At t hours after 10AM, the leaking truck is $50 \ln(t+2)$ miles from the garage, and the cleanup crew is $35t$ miles from the garage. You may use your calculator to evaluate any definite integrals for this problem.

a. [4 points] Calculate the total amount of paint that has leaked from the truck by 11AM.

b. [2 points] At time t hours after 10AM, what interval I of the road is still covered in paint? (you may assume that t represents a time before the trucks meet)

$$I = \left[\quad , \quad \right]$$

c. [3 points] Let $P(t)$ represent the amount of paint in gallons on the road t hours after 10 AM. Find a formula (which may include a definite integral) for $P(t)$.

d. [2 points] Calculate $P'(1)$.

8. [12 points] Sand dunes come in many shapes. *Barchan* dunes, which have the shape shown on the left, are studied extensively by geomorphologists. Horizontal cross-sections of these dunes are crescent-shaped (the dashed line encloses one such cross-section), and can be approximated as the shape on the right. The area of this shape is given by the formula $A_h = K\left(\frac{\pi}{2}Q_2 - \frac{4}{3}Q_1\right)$.



You are studying a barchan dune of 10 meters height, for which the values of Q_1 , Q_2 , and K vary with respect to the height h (in meters) of the cross-section according to the functions $Q_1(h) = 10 - h$, $Q_2(h) = 20 - 2h$, $K(h) = 100 - h^2$. The density of sand in the dune is $\delta = 1600$ kilograms per cubic meter.

- a. [5 points] Write an expression for the volume of one slice of sand dune h meters above the ground and Δh meters thick.
- b. [5 points] Write a definite integral that represents the total mass of sand in the dune. You do not need to evaluate this integral.
- c. [2 points] Write an expression (involving integrals) for the height of the center of mass of the sand dune. You do not need to evaluate this integral.