

# Math 116 — Second Midterm

March 20, 2013

Name: \_\_\_\_\_

Instructor: \_\_\_\_\_ Section: \_\_\_\_\_

1. **Do not open this exam until you are told to do so.**
2. This exam has 12 pages including this cover. There are 8 problems. Note that the problems are not of equal difficulty, so you may want to skip over and return to a problem on which you are stuck.
3. Do not separate the pages of this exam. If they do become separated, write your name on every page and point this out to your instructor when you hand in the exam.
4. Please read the instructions for each individual problem carefully. One of the skills being tested on this exam is your ability to interpret mathematical questions, so instructors will not answer questions about exam problems during the exam.
5. Show an appropriate amount of work (including appropriate explanation) for each problem, so that graders can see not only your answer but how you obtained it. Include units in your answer where that is appropriate.
6. You may use any calculator except a TI-92 (or other calculator with a full alphanumeric keypad). However, you must show work for any calculation which we have learned how to do in this course. You are also allowed two sides of a  $3'' \times 5''$  note card.
7. If you use graphs or tables to find an answer, be sure to include an explanation and sketch of the graph, and to write out the entries of the table that you use.
8. **Turn off all cell phones and pagers**, and remove all headphones.

---

Problem	Points	Score
1	12	
2	13	
3	10	
4	13	
5	12	
6	11	
7	15	
8	14	
Total	100	

1. [12 points] Indicate if each of the following is true or false by circling the correct answer. No justification is required.

a. [2 points] Consider the parametric equation given by  $x = a(1 + t^2)$  and  $y = 1 - t^3$ , where  $a > 0$ . Then the curve is concave up at the point  $(x, y) = (2a, 0)$ .

True                  False

b. [2 points] Let  $f(x)$  be a continuous function satisfying  $\lim_{x \rightarrow \infty} f(x) = 0$ . Then

$$\lim_{b \rightarrow \infty} \int_b^{\infty} f(x) dx = 0.$$

True                  False

c. [2 points] The point  $P$  whose polar coordinates  $(r, \theta) = (1, \frac{\pi}{6})$  also has coordinates  $(r, \theta) = (-1, \frac{7\pi}{6})$ .

True                  False

d. [2 points]  $\int_0^2 \ln(1 + t) dt$  is an improper integral.

True                  False

e. [2 points] All the solutions  $y(t)$  of the differential equation  $\frac{dy}{dt} = t^3$  are concave up.

True                  False

f. [2 points] The length of the parametric curve given by  $x = \cos t$  and  $y = \cos t + 1$  is  $2\sqrt{2}$ .

True                  False


2. [13 points]  
 a. [7 points] Consider the following differential equations:

A.  $y' = y(x - 2)^2$     B.  $y' = y(x - 2)$     C.  $y' = -y(1 - y)$     D.  $y' = -y^2(1 - y)$

Each of the following slope fields belongs to one of the differential equations listed above. Indicate which differential equation on the given line. Find the equation of the equilibrium solutions and their stability. If a slope field has no equilibrium solutions, write none.


Differential equation: \_\_\_\_\_

Equilibrium solutions and stability:




Differential equation: \_\_\_\_\_

Equilibrium solutions and stability:



- b. [4 points] A bank account earns a  $p$  percent annual interest compounded continuously. Continuous payments are made out of the account at a rate of  $q$  thousands of dollars per year. Let  $B(t)$  be the amount of money (**in thousands of dollars**) in the account  $t$  years after the account was opened. Write the differential equation satisfied by  $B(t)$ .
- c. [2 points] The slope field shown below corresponds to the differential equation satisfied by  $B(t)$  (for certain values of  $p$  and  $q$ ). Sketch on the slope field below the solution to the differential equation that corresponds to an account opened with an initial deposit of 3,000 dollars.



3. [10 points] The function  $y(t)$  satisfies the differential equation

$$\frac{dy}{dt} + 2y = 2t \quad \text{with} \quad y(0) = 1.$$

- a. [1 point] Can you use the method of separation of variables to solve this differential equation?
- b. [4 points] Use Euler's method with two steps to estimate  $y(1)$ . Fill the table with the values you find.

$t$	0		
$y(t)$			

- c. [5 points] For what values of  $a$  and  $b$  is the function  $y(t) = ae^{-2t} + b + t$  a solution to the differential equation

$$\frac{dy}{dt} + 2y = 2t \quad \text{with} \quad y(0) = 1 \quad a = \underline{\hspace{2cm}} \quad b = \underline{\hspace{2cm}}.$$


4. [13 points]

a. [8 points] Consider the functions  $f(x)$  and  $g(x)$  where

$$\frac{1}{x^2} \leq g(x) \quad \text{for} \quad 0 < x < \frac{1}{2}.$$

$$g(x) \leq \frac{1}{x^2} \quad \text{for} \quad 1 < x$$

$$\frac{1}{x^2} \leq f(x) \quad \text{for} \quad 1 < x.$$



Using the information about  $f(x)$  and  $g(x)$  provided above, determine which of the following integrals is convergent or divergent. Circle your answers. If there is not enough information given to determine the convergence or divergence of the integral circle NI.



- |                               |            |           |    |
|-------------------------------|------------|-----------|----|
| i) $\int_1^{\infty} f(x) dx$  | CONVERGENT | DIVERGENT | NI |
| ii) $\int_1^{\infty} g(x) dx$ | CONVERGENT | DIVERGENT | NI |
| iii) $\int_0^1 f(x) dx$       | CONVERGENT | DIVERGENT | NI |
| iv) $\int_0^1 g(x) dx$        | CONVERGENT | DIVERGENT | NI |

b. [5 points] Does  $\int_e^{\infty} \frac{1}{x(\ln x)^2} dx$  converge or diverge? If the integral converges, compute its value. Show all your work. Use u substitution.

5. [12 points] A particle moves according to the following parametric equations

$$x = x(t) \quad \text{and} \quad y = y(t) \quad \text{for} \quad -2 \leq t \leq 2,$$

where the graphs of  $x(t)$  and  $y(t)$  are shown below.



- a. [2 points] Is there a value of  $t$  at which the particle is at the point  $(0, 2)$ ? If so, find the value of  $t$  where this happens.
- b. [3 points] At which value(s) of  $t$  does the particle on the  $x$ -axis?
- c. [4 points] At what points  $(x, y)$  does the curve traveled by the particle have a horizontal tangent line? Include the time of each point.
- d. [3 points] For which of values of  $t$  is the slope of the tangent line to the curve positive?

6. [11 points]

- a. [8 points] Use the **comparison method** to determine the convergence or divergence of the following improper integrals. Justify your answers. Make sure to properly cite any results of convergence or divergence of integrals that you use.

i) 
$$\int_1^{\infty} \frac{3 + \sin(4x)}{\sqrt[3]{x}} dx.$$

ii) 
$$\int_4^{\infty} \frac{1}{\sqrt{x} + x^2} dx.$$

- b. [3 points] For which values of  $p$  does the following integral converges?

$$\int_2^{\infty} \frac{x^2 - 1}{x^p + 4x^2 + 2} dx.$$


No justification is required.



7. [15 points] A cone is filled with water up to a depth of  $H_0$  m. At time  $t = 0$ , a valve at the bottom of the cone is opened. Water leaks out of the cone through the opened valve. Let  $H(t)$  be the depth of the water (in m) in the cone at time  $t$  (in hours). The function  $H(t)$  satisfies the differential equation

$$\frac{dH}{dt} = \frac{k}{H^{\frac{3}{2}}}$$

- a. [2 points] What must be the sign and units of  $k$ ?



- b. [7 points] Find a formula for  $H(t)$ . Your formula should include  $k$  and  $H_0$

*This problem continues on the next page.*


***Problem 7 continued***

- c. [4 points] If the cone is filled with water up to a depth of 4 m at  $t = 0$ . What should the value of  $k$  be in order for the cone to be empty after an hour? Show all your work.

- d. [2 points] Does the differential equation satisfied by  $H$  have equilibrium solutions? If it does, find them.

8. [14 points]


- a. [6 points] Find a definite integral that computes the shaded area outside the circle  $r = \frac{5}{2}$  and inside the curve given by  $r = 2 + \sin 2\theta$  in the graph below.



- b. [4 points] Find the polar coordinates  $(r, \theta)$  of the points where the line  $y = \sqrt{3}x$  intersects the graph of  $r = 2 + b \sin 2\theta$ . Here the constant  $0 < b < 2$ . Your answers may include  $b$ .

$P_1 =$  \_\_\_\_\_

$P_2 =$  \_\_\_\_\_



c. [4 points]

i) Find the equation in polar coordinates of the line  $x = 0$ .

ii) Find the equation in polar coordinates of the line  $y = 4$ .