

4. (8 points) Show that if  $\sum_{n=1}^{\infty} a_n$  converges, then  $\lim_{n \rightarrow \infty} a_n = 0$ .

Let  $S_n = a_1 + a_2 + \cdots + a_n$ , so that  $S_n - S_{n-1} = a_n$ . The series converges to  $S$  if and only if  $S = \lim_{n \rightarrow \infty} S_n$  exists. In this case,  $\lim_{n \rightarrow \infty} S_{n-1} = S$  as well so

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} S_n - S_{n-1} = \lim_{n \rightarrow \infty} S_n - \lim_{n \rightarrow \infty} S_{n-1} = S - S = 0.$$

5. (8 points) In this question we will investigate the convergence of the power series  $\sum_{n=0}^{\infty} \frac{n^2}{e^n} (x+2)^n$ .

(a) Find the radius of convergence,  $R$ , of the power series. (Show your work.)

**Solution.** We will use the ratio test. Let  $a_n = \frac{n^2(x+2)^n}{e^n}$  so that

$$\frac{a_{n+1}}{a_n} = \frac{\frac{(n+1)^2(x+2)^{(n+1)}}{e^{n+1}}}{\frac{n^2(x+2)^n}{e^n}} = \frac{(n+1)^2(x+2)}{en^2}.$$

Consequently,

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{(n+1)^2}{en^2} |x+2| = \frac{|x+2|}{e}.$$

By the ratio test, the series therefore converges when this limit is less than one and diverges when it is greater than one. That is, the series converges for  $|x+2| < e$  and diverges for  $|x+2| > e$ , so the radius of convergence is  $R = e$ . The series converges for  $-e < x+2 < e$  or  $-2-e < x < -2+e$ .

$R = \underline{\quad \boxed{e} \quad}.$

(b) What is the interval of convergence of the power series?

$$\boxed{-2 - e < x < -2 + e.}$$