


1. [6 points] Suppose that the yield of the tomato plants in a particular Michigan garden is a function of the amount of water that the plants receive (from rainfall and irrigation). Let $T(w)$ be the seasonal yield (in pounds) of the tomato plants in a season when the plants receive w gallons of water every week. A portion of the graph of $T'(w)$ (the derivative of $T(w)$) is shown below.

Note that $T'(w)$ is linear for $50 \leq w \leq 70$. Let A be the area of the region between the w -axis and the graph of $T'(w)$ for $0 \leq w \leq 30$, and let B be the area of the region between the w -axis and the graph of $T'(w)$ for $30 \leq w \leq 50$,



- a. [2 points] If the tomato plants yield 150 pounds of tomatoes when the plants receive 70 gallons of water every week, how many pounds of tomatoes would the plants yield in a season when they receive 30 gallons of water each week? (Your answer may involve the constants A and B .)

Answer: _____

- b. [2 points] In order to maximize the yield of the tomato plants, how many gallons of water should the plants receive each week? (Round to the nearest 5 gallons.)

Answer: _____

- c. [2 points] Consider the integral $\int_{10}^{30} T'(w) dw$.

Rank the following four estimates of the value of this integral in order from least to greatest by writing them in the correct order on the answer blanks below:

LEFT(10) RIGHT(10) TRAP(10) MID(10)

_____ < _____ < _____ < _____