- **5.** [15 points] For each of the following, identify the statement as true or false by circling "True" or "False" as appropriate, and provide a short (one or two sentence) explanation indicating why that answer is correct.
  - **a**. [3 points] For the system  $x' = -xy + y^2$ ,  $y' = x^2 xy$ , the nonlinear trajectory in the phase plane with x(0) = -3 and y(0) = 0 lies on a circle centered on the origin.

True False

**b.** [3 points] For a linear differential operator  $L = \frac{d^2}{dt^2} + p(t)\frac{d}{dt} + q(t)$ , if  $y_1$  and  $y_2$  are different functions satisfying  $L[y_1] = L[y_2] = g(t) \neq 0$ , then, for any constants  $c_1$  and  $c_2$ ,  $y = c_1y_1 - c_2y_2$  satisfies L[y] = 0.

True False

c. [3 points] The solution to a differential equation my'' + ky = F(t) modeling the motion y of an undamped mechanical spring system with a periodic external force  $F(t) = F_0 \cos(\omega t)$  can always be written as  $y = A \cos(\omega_0 t - \delta_1) + B \cos(\omega t - \delta_2)$ , a sum of two oscillatory terms. (A, B,  $\omega_0$ ,  $\delta_1$  and  $\delta_2$  are constants.)

True False

d. [3 points] If  $\lambda^2 + p\lambda + q = 0$  is the characteristic equation of a constant-coefficient linear differential equation L[y] = g(t), then solving for  $Y(s) = \mathcal{L}\{y(t)\}$  will result in an expression involving a product of  $(s^2 + ps + q)^{-1}$  with other terms.

True False

e. [3 points] If  $f(t) \neq 0$  has Laplace transform  $\mathcal{L}{f(t)} = F(s)$  and  $g(t) = \begin{cases} f(t), & 0 < t < c \\ 0, & t \ge c \end{cases}$ , then  $\mathcal{L}{g(t)} = (1 - e^{-sc})F(s)$ . True False